



The only T2D therapy approved by the FDA to reduce the risk of end-stage kidney disease, doubling of serum creatinine, CV death, and hospitalization for heart failure in adults with T2D and diabetic nephropathy with albuminuria > 300 mg/day³

In adults who have T2D and diabetic nephropathy (ie, DKD) with albuminuria > 300 mg/day, INVOKANA[®] is the only SGLT2i indicated to slow the progression of DKD and reduce the risk of HHF³⁻⁶

See next page for a selection of ICD-10 codes and information about submitting prior authorizations ►

CV=cardiovascular; DKD=diabetic kidney disease; HHF=hospitalization for heart failure; SGLT2i=sodium-glucose co-transporter 2 inhibitor; T2D=type 2 diabetes.

* US population with T2D from the National Health and Nutrition Examination Survey 1999-2012. DKD based on either eGFR or albuminuria criteria defined by Kidney Disease: Improving Global Outcomes. The overall prevalence of DKD based on either eGFR (CKD-EPI) or albuminuria criteria was 43.5% (95% CI: 41.6, 45.4).

References: **1.** Bailey RA, Wang Y, Shu V, Rupnow MF. Chronic kidney disease in US adults with type 2 diabetes: an updated national estimate of prevalence based on Kidney Disease: Improving Global Outcomes (KDIGO) staging. *BMC Res Notes*. 2014;7:415. **2.** What is diabetic kidney disease? National Institute of Diabetes and Digestive and Kidney Diseases. <https://www.niddk.nih.gov/health-information/diabetes/overview/preventing-problems/diabetic-kidney-disease>. Accessed August 11, 2020. **3.** INVOKANA[®] [prescribing information]. Titusville, NJ: Janssen Pharmaceuticals, Inc. **4.** Jardiance[®] [prescribing information]. Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals, Inc. **5.** Farxiga[®] [prescribing information]. Wilmington, DE: AstraZeneca Pharmaceuticals LP. **6.** Steglatro[™] [prescribing information]. Whitehouse Station, NJ. Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc.

INDICATIONS

INVOKANA[®] (canagliflozin) is indicated:

- as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus
- to reduce the risk of major adverse cardiovascular events (cardiovascular death, nonfatal myocardial infarction, and nonfatal stroke) in adults with type 2 diabetes mellitus and established cardiovascular disease (CVD)
- to reduce the risk of end-stage kidney disease (ESKD), doubling of serum creatinine, cardiovascular (CV) death, and hospitalization for heart failure in adults with type 2 diabetes mellitus and diabetic nephropathy with albuminuria greater than 300 mg/day

Limitations of Use

INVOKANA[®] is not recommended in patients with type 1 diabetes mellitus. It may increase the risk of diabetic ketoacidosis in these patients.

INVOKANA[®] is not recommended for use to improve glycemic control in adults with type 2 diabetes mellitus with an eGFR less than 30 mL/min/1.73 m². INVOKANA[®] is likely to be ineffective in this setting based upon its mechanism of action.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

- Serious hypersensitivity reaction to INVOKANA[®], such as anaphylaxis or angioedema
- Patients on dialysis

Please read additional Important Safety Information on the following pages. Please read full Prescribing Information and Medication Guide for INVOKANA[®].

Learn more about INVOKANA[®] at www.INVOKANAhcp.com

DIABETIC KIDNEY DISEASE (DKD)

- E11.2:** Type 2 diabetes mellitus with kidney complications
- E11.21:** Type 2 diabetes mellitus with diabetic nephropathy
- E11.22:** Type 2 diabetes mellitus with diabetic chronic kidney disease
- E11:** Type 2 diabetes mellitus + **N18:** Chronic kidney disease (CKD)

TYPE 2 DIABETES (T2D)

- E11:** Type 2 diabetes mellitus

CARDIOVASCULAR DISEASE (CVD)

- E11.5:** Type 2 diabetes mellitus with circulatory complications
- E11.51:** Type 2 diabetes mellitus with diabetic peripheral angiopathy without gangrene
- E11:** Type 2 diabetes mellitus + **one of the following codes:**

Myocardial infarction

- I21.09:** ST elevation (STEMI) myocardial infarction involving other coronary artery of anterior wall

Coronary artery disease

- I24.8:** Other forms of acute ischemic heart disease
- I25.2:** Old myocardial infarction
- I25.5:** Ischemic cardiomyopathy
- I25.89:** Other forms of chronic ischemic heart disease
- I25.9:** Chronic ischemic heart disease, unspecified
- I25.10:** Atherosclerotic heart disease of native coronary artery without angina pectoris

Collected on September 3, 2020, and subject to change.

This document is presented for informational purposes only and is not intended to provide reimbursement or legal advice, nor does it promise or guarantee coverage, levels of reimbursement, payment, or charge. Similarly, all HCPCS codes are supplied for informational purposes only and represent no statement, promise, or guarantee by Janssen Pharmaceuticals, Inc., that these codes will be appropriate or that reimbursement will be made. It is not intended to increase or maximize reimbursement by any payer. Laws, regulations, and policies concerning reimbursement are complex and are updated frequently. While we have made an effort to be current as of the issue date of this document, the information may not be as current or comprehensive when you view it. We strongly recommend you consult the payer organization for its reimbursement policies.

Reference: 1. Centers for Medicare & Medicaid Services. 2020 ICD-10-CM: 2020 code descriptions in tabular order. <https://www.cms.gov/Medicare/Coding/ICD10/2020-ICD-10-CM>. Accessed September 3, 2020.

IMPORTANT SAFETY INFORMATION for INVOKANA®, continued

WARNINGS AND PRECAUTIONS

- **Lower-Limb Amputation:** An increased risk of lower-limb amputations associated with INVOKANA® use versus placebo was observed in CANVAS (5.9 vs 2.8 events per 1000 patient-years) and CANVAS-R (7.5 vs 4.2 events per 1000 patient-years), two randomized, placebo-controlled trials evaluating patients with type 2 diabetes who had either established cardiovascular disease or were at risk for cardiovascular disease. The risk of lower-limb amputations was observed at both the 100-mg and 300-mg once-daily dosage regimens.

Amputations of the toe and midfoot (99 out of 140 patients with amputations receiving INVOKANA® in the two trials) were the most frequent; however, amputations involving the leg, below and above the knee, were also observed (41 out of 140 patients with amputations receiving INVOKANA® in the two trials). Some patients had multiple amputations, some involving both lower limbs.

Please read full Prescribing Information and Medication Guide for INVOKANA®.

CARDIOVASCULAR DISEASE (CVD), continued

Stroke

- I60.9:** Nontraumatic subarachnoid hemorrhage, unspecified
- I61.9:** Nontraumatic intracerebral hemorrhage, unspecified
- I63.22:** Cerebral infarction due to unspecified occlusion or stenosis of basilar artery
- I63.139:** Cerebral infarction due to embolism of unspecified carotid artery
- I63.239:** Cerebral infarction due to unspecified occlusion or stenosis of unspecified carotid artery

Peripheral arterial disease

- I70.209:** Unspecified atherosclerosis of native arteries of extremities, unspecified extremity
- I70.399:** Other atherosclerosis of unspecified type of bypass graft(s) of the extremities, unspecified extremity
- I73.9:** Peripheral vascular disease, unspecified
- I74.2:** Embolism and thrombosis of arteries of the upper extremities
- I74.3:** Embolism and thrombosis of arteries of the lower extremities
- I74.5:** Embolism and thrombosis of iliac artery
- I75:** Atheroembolism

Heart failure

- I50:** Heart failure
- I50.1:** Left ventricular failure, unspecified
- I50.20:** Unspecified systolic (congestive) heart failure
- I50.30:** Unspecified diastolic (congestive) heart failure
- I50.40:** Unspecified combined systolic (congestive) and diastolic (congestive) heart failure

Sample prior authorization form

The form below is an example of a prior authorization form. Requests for prior authorization to prescribe INVOKANA® should be completed based on independent clinical judgment. Be sure to provide documentation from the patient's medical history to support your request.

1 MEDICATION
INVOKANA® (for appropriate patients).
Please see Indications in #4 below.

2 STRENGTH
INVOKANA® is available in two dose strengths (100 mg and 300 mg).¹
Choose the dose that is appropriate for your patient.

3 FREQUENCY
The recommended starting dose of INVOKANA® is 100 mg once daily, taken before the first meal of the day.¹ Dose can be increased to 300 mg once daily in patients tolerating INVOKANA® 100 mg once daily who have an eGFR of 60 mL/min/1.73 m² or greater and require additional glycemic control.¹

4 DIAGNOSIS
Insert your patient's diagnosis here.

INDICATIONS

INVOKANA® (canagliflozin) is indicated:

- as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus
- to reduce the risk of major adverse cardiovascular events (cardiovascular death, nonfatal myocardial infarction, and nonfatal stroke) in adults with type 2 diabetes mellitus and established cardiovascular disease (CVD)
- to reduce the risk of end-stage kidney disease (ESKD), doubling of serum creatinine, cardiovascular (CV) death, and hospitalization for heart failure in adults with type 2 diabetes mellitus and diabetic nephropathy with albuminuria greater than 300 mg/day

INVOKANA® is not recommended for the treatment of type 1 diabetes or to improve glycemic control in patients with type 2 diabetes and an eGFR less than 30 mL/min/1.73 m².

Formulary Exception/Prior Authorization Request Form

Patient Information				Prescriber Information			
Patient Name:				Prescriber Name:			
Patient ID#:				Address:			
Address:				City:			
City:		State:		City:		State:	
Home Phone:		ZIP:		Office Phone #:		Office Fax #:	
ZIP:		DOB:		Contact Person at Doctor's Office:			
Gender: M or F							
Diagnosis and Medical Information							
Medication:				Strength:		Directions for use (Frequency):	
Expected Length of Therapy:				City:		Day Supply:	
						If this is a continuation of therapy, how long has the patient been on the medication?	
Diagnosis:				Diagnosis (ICD) Code(s):			
FORM CANNOT BE EVALUATED WITHOUT REQUIRED CLINICAL INFORMATION							
<input type="checkbox"/> Expedited/Urgent Review Requested: By checking this box and signing below, I certify that applying the standard review time frame may seriously jeopardize the life or health of the patient or the patient's ability to regain maximum function.							
What condition is the drug being prescribed for? _____							
Please list all medications the patient has tried specific to the diagnosis and specify below: Medication name, reason for failure, including trial year: _____							
Drug(s) contraindicated: _____							
Adverse event (e.g., toxicity, allergy) for each drug: _____							

5 ICD-10 CODE(S)
Insert the appropriate ICD-10 code(s) for your patient on page 2. The ICD-10 codes provided here are for T2D, CVD, and DKD. Select any appropriate disease-specific code(s) based on your patient's diagnosis. There may be additional codes that are utilized for T2D, CVD, and DKD. For additional codes please refer to a coding resource or www.cms.gov.

6 PRIOR MEDICATIONS
Provide a brief account of your patient's prior medication history, including medication name, reason for failure, and trial year.

CVD=cardiovascular disease; DKD=diabetic kidney disease; T2D=type 2 diabetes.

Collected on September 3, 2020, and subject to change.

This document is presented for informational purposes only and is not intended to provide reimbursement or legal advice, nor does it promise or guarantee coverage, levels of reimbursement, payment, or charge. Similarly, all HCPCS codes are supplied for informational purposes only and represent no statement, promise, or guarantee by Janssen Pharmaceuticals, Inc., that these codes will be appropriate or that reimbursement will be made. It is not intended to increase or maximize reimbursement by any payer. Laws, regulations, and policies concerning reimbursement are complex and are updated frequently. While we have made an effort to be current as of the issue date of this document, the information may not be as current or comprehensive when you view it. We strongly recommend you consult the payer organization for its reimbursement policies.

Reference: 1. INVOKANA® [prescribing information]. Titusville, NJ: Janssen Pharmaceuticals, Inc.

IMPORTANT SAFETY INFORMATION for INVOKANA®, continued

WARNINGS AND PRECAUTIONS, continued

- **Lower-Limb Amputation (continued):** Lower-limb infections, gangrene, and diabetic foot ulcers were the most common precipitating medical events leading to the need for an amputation. The risk of amputation was highest in patients with a baseline history of prior amputation, peripheral vascular disease, and neuropathy.

Please read full [Prescribing Information](#) and [Medication Guide](#) for INVOKANA®.

IMPORTANT SAFETY INFORMATION for INVOKANA[®], continued

WARNINGS AND PRECAUTIONS, continued

- **Lower-Limb Amputation (continued):** Before initiating INVOKANA[®], consider factors in the patient history that may predispose to the need for amputations, such as a history of prior amputation, peripheral vascular disease, neuropathy, and diabetic foot ulcers. Counsel patients about the importance of routine preventative foot care. Monitor patients for signs and symptoms of infection (including osteomyelitis), new pain or tenderness, sores, or ulcers involving the lower limbs, and discontinue if these complications occur.
- **Volume Depletion:** INVOKANA[®] can cause intravascular volume contraction, which may sometimes manifest as symptomatic hypotension or acute transient changes in creatinine. There have been postmarketing reports of acute kidney injury which are likely related to volume depletion, some requiring hospitalizations and dialysis, in patients with type 2 diabetes mellitus receiving SGLT2 inhibitors, including INVOKANA[®]. Patients with impaired renal function (eGFR less than 60 mL/min/1.73 m²), elderly patients, or patients on loop diuretics may be at increased risk for volume depletion or hypotension. Before initiating INVOKANA[®] in patients with one or more of these characteristics, assess and correct volume status. Monitor for signs and symptoms of volume depletion after initiating therapy.
- **Ketoacidosis:** Ketoacidosis, a serious life-threatening condition requiring urgent hospitalization, has been identified in patients with type 1 and 2 diabetes mellitus receiving SGLT2 inhibitors, including INVOKANA[®]. In placebo-controlled trials of patients with type 1 diabetes, the risk of ketoacidosis was increased in patients who received SGLT2 inhibitors compared to patients who received placebo. The risk of ketoacidosis may be greater with higher doses. Fatal cases of ketoacidosis have been reported in patients taking INVOKANA[®]. Before initiating INVOKANA[®], consider factors in patient history that may predispose to ketoacidosis. For patients who undergo scheduled surgery, consider temporarily discontinuing INVOKANA[®] for at least 3 days prior to surgery. Monitor for ketoacidosis and temporarily discontinue in other clinical situations known to predispose to ketoacidosis. Ensure risk factors for ketoacidosis are resolved prior to restarting therapy. Educate patients on the signs and symptoms of ketoacidosis and instruct patients to discontinue INVOKANA[®] and seek medical attention immediately if signs and symptoms occur.
- **Urosepsis and Pyelonephritis:** Serious urinary tract infections, including urosepsis and pyelonephritis, requiring hospitalization have been reported in patients receiving SGLT2 inhibitors, including INVOKANA[®]. Treatment with SGLT2 inhibitors increases this risk. Evaluate for signs and symptoms and treat promptly.
- **Hypoglycemia With Concomitant Use With Insulin and Insulin Secretagogues:** Insulin and insulin secretagogues are known to cause hypoglycemia. INVOKANA[®] may increase the risk of hypoglycemia when combined with insulin or an insulin secretagogue. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with INVOKANA[®].
- **Necrotizing Fasciitis of the Perineum (Fournier's Gangrene):** Necrotizing fasciitis of the perineum, a rare but serious and life-threatening necrotizing infection requiring urgent surgical intervention, has been identified in postmarketing surveillance in female and male patients with diabetes mellitus receiving SGLT2 inhibitors, including INVOKANA[®]. Serious outcomes have included hospitalization, multiple surgeries, and death. If suspected, start treatment immediately with broad-spectrum antibiotics and, if necessary, surgical debridement. Discontinue INVOKANA[®].
- **Genital Mycotic Infections:** INVOKANA[®] increases risk of genital mycotic infections, especially in uncircumcised males or patients with prior infections. Monitor and treat appropriately.
- **Hypersensitivity Reactions:** Hypersensitivity reactions, including angioedema and anaphylaxis, were reported with INVOKANA[®]; these reactions generally occurred within hours to days after initiation. If reactions occur, discontinue INVOKANA[®], treat, and monitor until signs and symptoms resolve.
- **Bone Fracture:** Increased risk of bone fracture, occurring as early as 12 weeks after treatment initiation, was observed in patients using INVOKANA[®]. Prior to initiation, consider factors that contribute to fracture risk.

Please read full [Prescribing Information](#) and [Medication Guide](#) for INVOKANA[®].

IMPORTANT SAFETY INFORMATION for INVOKANA[®], continued

DRUG INTERACTIONS

- **UGT Enzyme Inducers:** Co-administration with rifampin lowered INVOKANA[®] exposure, which may reduce the efficacy of INVOKANA[®]. For patients with eGFR ≥ 60 mL/min/1.73 m², if an inducer of UGTs (eg, rifampin, phenytoin, phenobarbital, ritonavir) is co-administered with INVOKANA[®], increase the dose to 200 mg (taken as two 100 mg tablets) once daily in patients currently tolerating INVOKANA[®] 100 mg. The dose may be increased to 300 mg once daily in patients currently tolerating INVOKANA[®] 200 mg and who require additional glycemic control.

For patients with eGFR < 60 mL/min/1.73 m², if an inducer of UGTs is co-administered with INVOKANA[®], increase the dose to 200 mg (taken as two 100 mg tablets) once daily in patients currently tolerating INVOKANA[®] 100 mg. Consider adding another antihyperglycemic agent in patients who require additional glycemic control.

- **Digoxin:** There was an increase in the AUC and mean peak drug concentration of digoxin when co-administered with INVOKANA[®] 300 mg. Monitor appropriately.
- **Positive Urine Glucose Test:** Monitoring glycemic control with urine glucose tests is not recommended in patients taking SGLT2 inhibitors. Use alternative methods to monitor glycemic control.
- **Interference With 1,5-Anhydroglucitol (1,5-AG) Assay:** Monitoring glycemic control with 1,5-AG assay is not recommended in patients taking SGLT2 inhibitors. Use alternative methods to monitor glycemic control.

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** INVOKANA[®] is not recommended in pregnant women, especially during the second and third trimesters.
- **Lactation:** INVOKANA[®] is not recommended while breastfeeding.
- **Pediatric Use:** Safety and effectiveness in patients < 18 years of age have not been established.
- **Geriatric Use:** Patients ≥ 65 years had a higher incidence of adverse reactions related to reduced intravascular volume, particularly with the 300-mg dose; more prominent increase in the incidence was seen in patients who were ≥ 75 years. Smaller reductions in HbA1c relative to placebo were seen in patients ≥ 65 years.
- **Renal Impairment:** The efficacy and safety of INVOKANA[®] for glycemic control were evaluated in a trial that included patients with moderate renal impairment (eGFR 30 to < 50 mL/min/1.73 m²). These patients had less overall glycemic efficacy, and patients treated with 300 mg per day had increases in serum potassium, which were transient and similar by the end of the study. Patients with renal impairment using INVOKANA[®] for glycemic control may be more likely to experience hypotension and may be at a higher risk for acute kidney injury. INVOKANA[®] is contraindicated in patients with ESKD on dialysis.
- **Hepatic Impairment:** INVOKANA[®] has not been studied in patients with severe hepatic impairment and is not recommended in this population.

OVERDOSAGE

- In the event of an overdose, contact the Poison Control Center and employ the usual supportive measures.

ADVERSE REACTIONS

- The most common adverse reactions associated with INVOKANA[®] (5% or greater incidence) were female genital mycotic infections, urinary tract infections, and increased urination.

Please read accompanying full Prescribing Information and Medication Guide for INVOKANA[®].

Need more access & support info?

Visit www.INVOKANAhcp.com

Janssen Pharmaceuticals, Inc.

Canagliflozin is licensed from Mitsubishi Tanabe Pharma Corporation.

© Janssen Pharmaceuticals, Inc. 2020 September 2020 cp-109421v4

